

Heavy quark distribution function in hadrons

A.G.Oganesian

Institute of Theoretical and Experimental Physics, Moscow, Russia

Abstract

The moments of the heavy quark-parton distribution functions in a heavy pseudoscalar meson, obtained in QCD sum rules, are expanded in the inverse heavy quark. Comparison with the finite mass results reveals that while the heavy mass expansion works reasonably well for the b quark, one has to take into account terms of higher than $(1/m_c)^2$ order for the c quark.

In this talk I will very shortly report the result, obtained recently in my paper [1]. The problem is that there exist a large number of different theoretical models and approaches usually used to describe the heavy quark fragmentation functions (FF)(see, for example [2 – 6]), and expansions, based on the heavy-quark mass limit or on HQET are widely used in many of them (e.g. [6]). So the question arise, does the expansion work good, especially for charm case. In [1] it was offered to check the applicability (and accuracy) of this expansion using the moments of the heavy-quark structure obtained long ago from QCD sum rules in [7]. In this approach the moments of the heavy-quark parton distribution functions $M_n^c = \int_0^1 dx x^{n-1} c(x, Q^2)$ were directly calculated in QCD sum rules and FF were estimated, following the familiar relation [8] (see also [2] and [7] for details). The advantage of this method is the possibility to estimate the heavy parton distribution functions (hence, also fragmentation functions) in full QCD, in terms of universal parameters, such as quark masses and condensate densities. One can use this result to fix the parameters of different models, as was discussed in [7].

There is no time to discuss the method itself, one can look it in the paper [7] and also [1]. So I very briefly remind the main points of the method. One starts from the four-point correlator:

$$\Pi_{\mu\nu} = i \int e^{ip_1x+iq_1y-ip_2z} d^4x d^4y d^4z \langle 0 | T \{ j_5(x) j_\mu^{\text{em}}(y) j_\nu^{\text{em}}(0) j_5^\dagger(z) \} | 0 \rangle, \quad (1)$$

where $j_\mu^{\text{em}} = \bar{c} \Gamma_\mu c$, and heavy-light pseudoscalar currents $j_5 = \bar{c} \Gamma_5 u$ interpolate D meson. The above correlator is considered in the deep spacelike region at $t = (p_1 - p_2)^2 = 0$, $p_i^2, q_i^2 < 0$. We take into account the contributions of the unit operator (bare loop), quark condensate $a = -(2\pi)^4 \langle 0 | \bar{\psi} \psi | 0 \rangle$ and quark–gluon condensate $\langle 0 | \bar{\psi} G_{\mu\nu} \sigma^{\mu\nu} \psi | 0 \rangle$. The gluon and four-quark condensate contributions are neglected, because the estimates show, that they are very small. We express this amplitude in terms of double dispersion

relation over p_1^2 and p_2^2 , then, us usual, saturate the amplitude by hadronic states and equate it to the result of OPE and perform two independent Borel transformations on p_1^2 and p_2^2 . (We will equate the Borel masses $M_{1B}^2 = M_{2B}^2 = M_B^2$ only in the final sum rule). Following [7], we choose the value of the variable $s = (p_1 + q_1)^2$ in the unphysical region as $s = m_D^2 - Q^2$, and transform the integrals into the moments of the parton distribution. Finally the result, obtained in, is [7]:

$$M_n = g_D^{-2} m_D^{-4} \left(\int_{m^2}^{s_0} du_1 \int_{z_-}^{z_+} dz \frac{f_0 e^{-2(u_1 - m_D^2)/M_B^2}}{(1 + z - m_D^2/Q^2)^{n+1} \pi^2} + \frac{maL e^{-2(m^2 - m_D^2)/M_B^2}}{4\pi^2 \eta^{n+1}} + R_6 \right) \quad (2)$$

Here,

$$f_0 = \frac{3}{8} (s_1 - m^2) \left[\frac{1 + z + (s_1 + 2m^2)/Q^2}{((1 + z - s_1/Q^2)^2 + 4s_1/Q^2)^{-1/2}} - \frac{4m^2}{Q^2} \left(1 + \frac{4m^2}{Q^2} \right)^{-1/2} \right] \quad (3)$$

and

$$R_6 = \frac{mm_0^2 a}{4\pi^2} \frac{1}{2M_B^2} \left[4/M_B^2 - 2m^2/M_B^4 + 1/Q^2 + \frac{(n+1)}{\eta} \left(\frac{2}{3Q^2} - 1/M_B^2 - \frac{2m^2}{M_B^2 Q^2} \right) - \frac{(n+1)(n+2)m^2}{Q^4 \eta^2} \right] \eta^{-n-1} e^{-2(m^2 - m_D^2)/M_B^2}, \quad (4)$$

Here $\eta = (1 - (m^2 - m_D^2)/Q^2)$. The factor L accounts for the anomalous dimension of the quark condensate, normalized at point μ . We adopt the values of condensates [9, 10], normalizing them at $\mu = 1$ GeV. In this case the factor L is close to the unity.

In [7], it was shown that the first few moments of the sum rules are well behaved and the following numerical results were obtained at $Q_0^2 = 20$ GeV² ($Q_0^2 = 10$ GeV²): $M_2 = 0.85(0.9)$, $M_3 = 0.75(0.83)$, $M_4 = 0.67(0.78)$. These moments can be used as an input in the evolution equations, at some initial point $Q_0^2 = (3 - 5)m_D^2$, hence it is possible to predict the structure functions also at large Q^2 .

Now we can answer to our question if it is possible to use the heavy-quark limit for the heavy quark parton distribution (and, correspondingly, fragmentation function). For that, we expand the sum rules for the moments in the inverse mass of the heavy quark.

Substituting in Eq.(4) instead of the $g_D^2 m_D^4$ the sum rule obtained from the two-point correlator [11] we can rewrite the sum rules for M_n as

$$M_n = \frac{R_0(n) + R_4(n) + R_6(n)}{K_0 + K_4 + K_6}, \quad (5)$$

where the terms $R_d(n)$ in the numerator and K_d in the denominator originate from the dimension $d = 0, 4, 6$ contributions to the OPE of the four-point and two-point correlator, respectively. Following [7] we adopt the relation $M_B^2/2 = M^2$ between the Borel masses in the four-point and two-point sum rules. Note that the exponential factor containing m_D^2 cancels in the ratio of the two sum rules.

To investigate the heavy-quark mass limit of (5) we employ the standard scaling relations for the heavy hadron mass, continuum threshold, and Borel parameter: $m_D = m + \bar{\Lambda}$, $s_0 = m^2 + 2m\omega$, $M^2 = 2\tau m$ and in addition assume the scaling $Q^2 = \gamma m_D^2$, where in the heavy-quark limit the parameters $\bar{\Lambda}$, ω , τ and γ do not depend of the heavy quark mass. (One should note, that $\bar{\Lambda}$ here has nothing common with well-known QCD parameter Λ_{QCD}). In what follows we will use notation $\delta = \bar{\Lambda}/m$, $y = \bar{\Lambda}/\tau$.

Expanding in powers of $1/m$, we obtain the first three terms for R_0 :

$$R_0(n) \simeq D_2 + (n+1)\delta \frac{2D_2 - (2+\gamma)D_3}{\gamma} + \delta^2(n+1)(n+2) \left[\frac{2(n+2) + 3\gamma}{\gamma^2(n+2)} D_2 - \frac{2}{\gamma} \left(1 + \frac{2(n+1)}{(n+2)\gamma} \right) D_3 + \frac{2D_4}{3\gamma^2} \left(\gamma^2 + 4\gamma + 3 + \frac{2\gamma}{(n+1)(n+2)} \right) \right]. \quad (6)$$

Here, D_n is defined as $D_n = E_n(\omega/\tau)/y^{n-2}$ and $E_n(z) = \int_0^z dz e^{-z} z^n$. For the contribution of $d = 4$ operators we obtain:

$$R_4(n) \simeq \frac{a}{12\tau^3} L \left(1 + 2\delta(n+1)/\gamma + \delta^2 \frac{(n+1)(n+2)}{\gamma} [2/\gamma - 3/(n+2)] \right). \quad (7)$$

Finally, the contribution of $d = 6$ operators transforms into (first three terms)

$$R_6(n) \simeq \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} \left[-1 + \frac{2\delta}{y} \left(4 - (n+1)[1 + (y+2)/\gamma] \right) - \frac{2\delta^2(n+1)}{y\gamma} \left((n+2)[2 + (y+4)/\gamma + 2/(y\gamma)] - 3y - 24 - (8/3)(2n+5)/y \right) \right]. \quad (8)$$

And, finally, for the denominator $K = K_0 + K_4 + K_6$ in (5) we obtain after expansion

$$K \simeq D_2 - 2D_3\delta - 4D_4\delta^2 + \frac{a}{12\tau^3} \left(L - \frac{m_0^2}{16\tau^2} \right) + \frac{a}{12\tau^3} \frac{m_0^2}{16\tau^2} \frac{4\delta}{y}. \quad (9)$$

From these equations it is easy to calculate the actual value of moments in the first and second orders of expansion and compare it with the exact value to examine the accuracy of the expansion in the inverse heavy quark mass. For numerical analysis we choose $\bar{\Lambda} = 0.6$ GeV, $\omega = 1.4$ GeV, and $\tau = 0.6$ GeV, as usual. The numerical results for heavy-quark mass expansion of the first is shown in Fig. 1 where it is plotted as a function of the dimensionless ratio $k = m/m_c$, for two different values of the ratio $\gamma = Q^2/M_D^2 = 2, 4$. One can easily see, that even at relatively large masses of order of the b -quark mass ($k \sim 3$) the heavy mass limit and even the first-order expansion in inverse mass are not reliable. Including the second $O(1/m^2)$ term one improves the situation for the b -quark, but the approximate result for the moments still remains very far from the exact answer in the case of c -quark. We conclude that the heavy quark limit is not a reliable approximation for the parton distributions and fragmentation functions of c -quark. The analysis for higher moments (second and third) totally confirm this conclusion.




Figure 1: The first moment as a function of the quark mass m in the units $k = m/m_c$. Curves 1,2,3 correspond to the first-and second-order expansion in the inverse quark mass and to the exact answer, respectively. The value of $\gamma = Q^2/M_D^2 = 2(4)$ is chosen in the left (right) panel.

The author thanks B.L. Ioffe and A. Khodjamirian for useful discussions. This work was supported in part by the Russian Foundation of Basic Research, project no. 06-02-16905a and the funds from EC to the project "Study of the Strong Interacting Matter" under contract N0. R113-CT-2004-506078.

References

- [1] A. G. Oganesian, arXiv:0711.2830, to be published in Sov. J. Nucl. Phys.
- [2] V. G. Kartvelishvili, A. K. Likhoded, and V. A. Petrov, Phys. Lett.**B78** (1978) 615.
- [3] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, Phys. Rev.**D27**, (1983) 105.
- [4] M. Suzuki, Phys. Rev.**D33**, (1986) 676 .
- [5] P. D. B. Collins and T. P. Spiller, J.Phys.**G11**, (1985) 1289.
- [6] E. Braaten, K. M. Cheung, S. Fleming, and T. C. Yuan, Phys. Rev.**D51**, (1995) 4819.
- [7] G. L. Balaian, A. Y. Khodjamirian, and A. G. Oganesian, Sov.J.Nucl.Phys.**49**, (1989) 697
- [8] V. N. Gribov and L. N. Lipatov, Sov. J. Nucl. Phys. **15**, (1972) 675
- [9] B. V. Geshkenbein, B. L. Ioffe, and K. N. Zyablyuk, Phys. Rev.**D64**, 093009 (2001)

- [10] B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, (2006) 232
- [11] T. M. Aliev and V. L. Eletsky, Sov. J. Nucl.Phys. **38**, (1983) 936